

Field Crops

Crop	Dose per hectare	Seed dressing	Application 1	Application 2
Soybean (option A: 1 spray per season)	100mls per hectare	N/A	Spray 100mls per hectare at the 4 th to 6 th trifoliolate. Stage V4 – V6.	N/A
Soybean (option B: 2 sprays per season)		N/A	Spray 100mls per hectare at the 4 th to 6 th trifoliolate. Stage V4 – V6.	Spray 100mls per hectare at flowering. Stages R1 – R2.
Wheat (option A: 1 spray per season)		Mix 100mls of CropBioLife per 100 kgs of seed (optional)	Spray 100mls per hectare at Feekes 1-4	N/A
Wheat (option B: 2 sprays per season)		Mix 100mls of CropBioLife per 100 kgs of seed (optional)	Spray 100mls per hectare at Feekes 1-4	Spray 100mls per hectare at Feekes 5-7
Corn (option A: 1 spray per season)		N/A	Spray 100mls per hectare at stage V4 – V6	N/A
Corn (option B: 2 sprays per season)		N/A	Spray 100mls per hectare stage V4 – V6	Spray 100mls per hectare when tassel emerges (VT)
Alfalfa		N/A	Spray 100mls per hectare at 4 leaf stage	Spray 100mls per hectare at full flower
Sorghum or Milo		N/A	Spray 100mls per hectare at 4 leaf stage	Spray 100mls per hectare at full flower
Oats		Do not treat	Spray 100mls per hectare when 4 leaves have unfolded	Spray 100mls per hectare at full flower
Clover		N/A	Spray 100mls per hectare Spray about 1 week after clover sprout	N/A
Canola		Mix 100mls of CropBioLife per 100 kgs of seed (optional)	Spray 200mls per hectare at 5-10 leaf stage	Spray 150mls per hectare at Bud development

Orchards

Crop	Dose per hectare	Application 1	Application 2	Application 3
Apples	Spray 550mls per hectare	80-100% petal drop. The timing of this first application is critical to ensure best results	28 days later	28 days later
Almonds				
Cherries				
Citrus				

Ground fruit and vegetables

Crop	Dose per hectare	Seed dressing or plant dip	Application 1	Application 2	Additional
Tomatoes	300mls – 500mls per Hectare	(OPTIONAL) Dip small plants in a CropBioLife solution (50mls of CropBioLife per 100 litres of water)	Spray at the beginning of the season or 2 weeks after transplant	Spray 28 days later	Continue to spray every 28 days until harvest
Peppers					
Strawberries					
Blueberries					
Melons					
Zucchini					
Squash					
Cucumber					
Cantaloupe					
All other ground fruit or vegetables					

Wine grapes

Crop	Dose per hectare	Application 1	Application 2	Application 3
Wine grapes	300mls per hectare	Shoots 7 – 10 inches	Cap fall	Pea-sized berries

Table grapes

Crop	Dose per hectare	Application 1	Application 2	Application 3	Application 4
Table grapes	300mls per hectare	7-10 inch shoot size	Pre-flowering (preferable)	Flowering	Pea sized berries

Pasture, turf grass, home garden, lawns

Crop	Concentration in water	Application 1	Application 2	Application 3
Pasture, turf grass, home garden, lawns	Mix 20mls or 1 teaspoon of CropBioLife per 10 litres of water	Beginning of season	28 days later	28 days later

